

## Internet Addiction and Its Determinants among Medical Students in Dhaka City during COVID-19 Pandemic

Nusrat Jahan \*, Md. Rumman Idris, Iffat Tania Prianka, Pradip Kumar Sen Gupta

\*Department of Biochemistry, Ibrahim Medical College, Bangladesh

Cardiovascular Surgery Center, Combined Medical Hospital Dhaka, Bangladesh

Department of Community Medicine, Ibrahim Medical College, Bangladesh

Department of Epidemiology, Bangladesh University of Health Sciences, Bangladesh

### Abstract

#### Keywords:

*Internet Addiction, medical students, Covid-19, social distance.*

Internet addiction has now emerged as a global public health concern affecting people all over the world. This problem has been exacerbated by the COVID-19 epidemic, which has the potential to modify many social habits in young people. An online cross-sectional study of 439 medical students from 20 public and private medical colleges in Dhaka city was done using a semi-structured questionnaire and Young's internet addiction test. The study's goal was to assess the proportion of medical students who were identified to be addicted to the internet during the COVID-19 pandemic. Staying in private accommodation, spending more than 4 hours/day online, monthly internet expenditure, usual logging status, physical inactivity, and maintaining social distancing were found to be substantially associated factors with internet addiction among the sample. Taking the findings into account, it is critical to provide preventive measures and strengthen education on Internet Addiction, as well as to consolidate and promote mental health literacy, psychological warning signs, and adaptive psychiatric services during this trying time.

### Introduction

Internet Addiction (IA), by several sources, has been classified as a psychological reliance on the internet irrespective of the nature of activity engaged to which can also be referred to as an individual's inability to regulate their internet use on their own, later impacting their functionality (Bates et al., 2020; Nath, Naskar & Victor, 2016; Ghosh & Chatterjee, 2018). In 1995, Ivan Goldberg coined the expression "Internet Addiction" for which he suggested a medical criterion marked by (i) Maladaptive Internet Usage that causes pain or impairment in psychological, educational, or other essential functions, (ii) Tolerance, defined as the need to extend the duration of an internet connection to achieve the desired level of enthusiasm, and (iii) Withdrawal, defined as psychomotor restlessness or anxiety following the discontinuation or reduction of long-term Internet use. (Nath, et al., 2016; Ghosh & Chatterjee, 2018; Poli & Agrimi, 2012).

Excessive internet usage is reported to create morphological mutations of brain structure which has been reported in many studies. One of the studies found students using 10 hours in front of a computer for 6 days a week possess a smaller dorsolateral prefrontal cortex and rostral anterior cingulate cortex along with other parts of the cerebellum when compared with those of the control group (Uddin et al., 2016). Moreover, addicted people suffer from sleep disturbances, backaches, headaches, Carpal Tunnel Syndrome, blurred vision, dry eyes, and bad eating habits. Excessive use of Internet and addiction-like most other addiction, shelter one from dealing with difficult emotions and make them feel safer when indulged into and change their temporal experience (Salicetia, 2015). There has been found a strong association of Internet Addiction with conditions like depression, anxiety, fatigue, isolation, self-

esteem, social connectedness, insomnia, etc (Haand & Shuwang, 2020). Not only this, internet addiction makes people more susceptible to other addictions like drugs, alcohol, tobacco, sex, etc (Uddin et al., 2016).

Globally around 4.57 billion people, which is approximately 59% of the world population, are active internet users until July 2020 while the proportion of adolescent internet users has increased tremendously (Weinstein & Lejoyeux, 2010; Uddin et al., 2016). Like the rest of the world, the no. of internet users is also rising in Bangladesh and the current government is also promoting internet access through the "Digital Bangladesh" movement in the country (Mamun et al., 2019). From 2000 to 2016, the number of internet consumers increased in Bangladesh from 0.1 million to 62 million (Islam & Hossin, 2016).

As the world's use of the Internet grows, Internet Addiction has emerged as a new public health issue and globally Internet Addiction prevalence was found to be ranged from 3 percent to 38 percent in previous surveys (Chaudhari, Menon, Saldanha, Tewari & Bhattacharya, 2015). Similarly, Internet Addiction is gradually becoming a newly emerging problem for Bangladesh with the rise of internet users (Hassan, Alam, Wahab & Hawladar, 2020). In numerous Bangladeshi samples utilizing the Internet Addiction Test (IAT), prevalence rates of IA have been observed to range between 4 percent and 49.7 percent, which is comparable to other Asian countries (Hassan et al., 2020; Karim & Nigar, 2014, Uddin et al., 2016; Islam & Hossin, 2016; Mamun & Griffiths, 2019). The prevalence of Internet Addiction was found 27.1 percent in Hassan et al. (2020) whereas 34.3 percent of the Karim & Nigar (2014) study sample was reported to be moderate Internet users with 1.7 percent as heavy users.

Since the onset of the COVID-19 pandemic people have become used to operating their daily lives despite maintaining social distancing (Hall, Laddu, Phillips, Lavie & Arena, 2020). Adolescents are especially affected immensely by the closure of the learning institution, as well as other socio-behavioral changes like social distancing, lockdown, online class, etc. (Bates et al., 2020; Garcia-Priego, 2020). In an attempt to contain the pandemic's spread in Bangladesh, the government also issued a countrywide lockdown. As an aftermath, students have been seen to spend a significant amount of time online for academic, leisure, social needs and so on, which, if left unchecked, can lead to Internet Addiction (Dong, Yang, Lu & Hao, 2020).

Currently there is no research in Bangladesh that we are aware of that has looked into the pattern of Internet Addiction among medical students or how the COVID-19 pandemic may exacerbate this problem. During this crisis, assessing Internet Addiction level of medical students is as vital as the identification of the determinants that contribute to it.

## Materials and methods

### Subheading (10pt Times New Roman, Bold, Justified)

This online-based cross-sectional study was conducted on medical students of 20 different public and private medical colleges in Dhaka city over a period from August 2020 to January 2021. Total 439 sample were selected by using convenient sampling technique. Self-administrated semi-structured online questionnaire was used for the information regarding demographic and internet use variable. The online survey was conducted by using Google Forms, which were distributed through email and social media platforms. Regarding Internet Addiction, Young's 20-item Addiction test was used which is graded on a 5-point Likert scale to determine how much Internet use impacts a person's social life, sleep, academic activities, interpersonal connections, and emotions. The test's 20 questions are graded on a scale of 1 to 5, with 1 being the least severe and 5 being the most severe. The overall score indicates the degree of addiction and gives details on the negative consequences of Internet Addiction in numerous areas of life. Score interpretation:

1. **0 - 20:** are indicate a normal level of internet usage.
2. **21 - 49:** reflect the presence of a mild level of Internet Addiction.
3. **50 - 79:** reflect the presence of a moderate level of Internet Addiction.
4. **80 - 100:** indicate a severe level of Internet Addiction.
- 5.

Since the moderate users are often unable to control their internet use, we considered both moderate and excessive use of Internet (IAT total score  $\geq 50$ ) to be considered as IA (Internet Addicted), and who scored between 0 and 49 was considered as a normal user of Internet (Ali, Mohammad & Aly, 2017; Hassan et al., 2020; Islam & Hossin, 2016).

All statistical analysis were done by using the data analysis software SPSS 25.0. The data was summarized using descriptive statistics (frequency, percentage, and mean). To access the degree of association between the dependent and independent variable, Chi-square was applied. Multiple logistic regression analysis was used to determine the strength of association for variables that showed significant relationship in chi-square analysis. In all cases, P values of less than 0.05 were considered significant.

## Results

The sample was of a total of 439 medical students who were studying at different levels of the academic year of undergraduate medical courses. Majority of the students, 238 (54.2%), were in the age range 18-20 years and 299 students were female (68.1%). 307 (69.9%) were living with family and 308 (70.2%) respondents reported to live in urban area.

Most of the students were using the Internet for more than 5 years ( $n = 195$ , 44.4%). Among the students, 290 (66.1%) reported to be on the Internet more than 4 hours every day, and the most popular way of using the Internet was Wi-Fi 320 (72.9%). Majority of the students spent less than 800 taka per month on the internet ( $n = 245$ , 55.8%), and most common gadget was mobile ( $n = 418$ , 95.2%). Of the respondents, 50.3% said that they were occasionally online and Social media (55.6%) was the most accessed content, followed by academic materials (30.3%). Due to the COVID-19 pandemic, 266 (60.6%) students spent more than 5 hours weekly on online classes and 92.1% of students believed that their internet usage has increased due to maintaining social distancing. Among the students, 247 (56.3%) did not meet up the recommended physical activity; however, 192 (43.7%) students reported to meet the bar.




Figure 1: Distribution of the studied medical students according to Internet Addiction level

We considered moderate and severe Internet Addiction as Internet addict (score 50-100), which was 34.9%, and those with lower scores (0-49) were considered as non-Internet Addiction.

**Table 1. Association of demographic variables with Internet Addiction**

Variables	n	Addicted (%)	Non- addicted (%)	$\chi^2$ value	P-value
<b>Age</b>					
18-20 years	238	80(33.6%)	158(66.4%)	0.3561	0.553 <sup>ns</sup>
21-24 years	201	73(36.3%)	128(63.7%)		
<b>Gender</b>					
Male	140	54(38.6%)	86(61.4%)	1.252	0.263 <sup>ns</sup>
Female	299	99(33.1%)	200(66.9%)		
<b>Father's occupation</b>					
Govt. Employee	80	26(32.5%)	54(67.5%)	2.524	0.773 <sup>ns</sup>
Businessman	108	36(33.3%)	72(66.7%)		
Doctor	45	16(35.6%)	29(64.4%)		
Retired	27	13(48.1%)	14(51.9%)		
Others	139	49(35.3%)	90(64.7%)		
Teacher	40	13(32.5%)	27(67.5%)		
<b>Mother's occupation</b>					
Housewife	307	108(35.2%)	199(64.8%)	0.989	0.911 <sup>ns</sup>
Doctor	24	9(37.5%)	15(62.5%)		
Teacher	66	22(33.3%)	44(66.7%)		
Govt. Employee	16	4(25.0%)	12(75.0%)		
Others	26	10(38.5%)	16(61.5%)		
<b>Academic year</b>					
1 <sup>st</sup> year	201	73(36.3%)	128(63.7%)	0.351	0.553 <sup>ns</sup>
2 <sup>nd</sup> year	138	38(27.5%)	100(72.5%)	4.744	0.029*
3 <sup>rd</sup> year	43	14(32.6%)	29(67.4%)	0.111	0.739 <sup>ns</sup>
4 <sup>th</sup> year	34	15(44.1%)	19(55.9%)	1.393	0.237 <sup>ns</sup>
5 <sup>th</sup> year	23	13(56.5%)	10(43.5%)	5.020	0.025*
<b>Residence</b>					
Urban	308	107(34.7%)	201(65.3%)	0.006	0.997 <sup>ns</sup>
Semi-urban	74	26(35.1%)	48(64.9%)		
Rural	57	20(35.1%)	37(64.9%)		
<b>Living setup</b>					
Live with family	307	79(25.7%)	228(74.3%)	37.39	<0.001*
Live with other	132	74(56.1%)	58(43.9%)		

*Chi-squared Test ( 2) was done to analyze the data. \*significant*

Table 1 Shows Internet Addiction was significantly associated with living setup (P <0.001\*), students who were living with other people where more addicted to Internet than those who were living with family. The Association with Internet Addiction was found to be significant with students being in 2nd (P =0.029\*) and 5th year (P =0.025\*).

**Table 2. Association of Internet use and COVID-19 related variables with Internet Addiction**

Variables	N	Addicted (%)	Non- addicted (%)	$\chi^2$ value	P-value
<b>Length of internet uses (years)</b>					
More than 5	195	69(35.4%)	126(64.6%)	0.566	0.754 <sup>ns</sup>
2-5	179	64(35.8%)	115(64.2%)		
Less than 2	65	20(30.8%)	45(69.2%)		
<b>Daily time spent on Internet (hours)</b>					

More than 4	290	124(42.8%)	166(57.2%)	25.26	<0.001*
2-4	127	22(17.3%)	105(82.7%)		
Less than 2	22	7(31.8%)	15(68.2%)		
<b>The most common way to using the Internet</b>					
Wi-Fi	320	112(35.0%)	208(65.0%)	0.011	0.915 <sup>ns</sup>
Mobile data	119	41(34.5%)	78(65.5%)		
<b>Most used gadget</b>					
Mobile	418	142(34.0%)	276(66.0%)	2.984	0.084 <sup>ns</sup>
Personal desktop or laptop	21	11(52.4%)	10(47.6%)		
<b>Monthly expenditure for internet access (Taka)</b>					
800 or more	206	88(42.7%)	118(57.3%)	10.87	0.001*
Less than 800	233	65(27.9%)	168(72.1%)		
<b>Usual logging status</b>					
Always online	168	83(49.4%)	85(50.6%)	25.90	<0.001*
Occasionally check-in	221	55(24.9%)	166(75.1%)		
Infrequently check-in	46	14(30.4%)	32(69.6%)		
Deactivated or never check-in	4	1(25.0%)	3(75.0%)		
<b>Most accessed content on the Internet</b>					
Social media	244	92(37.7%)	152(62.3%)	6.070	0.194 <sup>ns</sup>
Academic materials	133	42(31.6%)	91(68.4%)		
Downloadable media	40	10(25.0%)	30(75.0%)		
Online shopping	4	3(75.0%)	1(25.0%)		
Others	18	6(33.3%)	12(66.7%)		
<b>Duration of online classes (hours/week)</b>					
More than 5	266	90(33.8%)	176(66.2%)	1.211	0.546 <sup>ns</sup>
2-5	128	44(34.4%)	84(65.6%)		
Less than 2	45	19(42.2%)	26(57.8%)		
<b>Meets recommended PA</b>					
No	247	103(41.7%)	144(58.3%)	11.666	<0.001*
Yes	192	50(26.0%)	142(74.0%)		
<b>Believe in the past few months internet usage has increased due to maintaining social distancing</b>					
No	35	5(14.3%)	30(85.7%)	7.085	0.008*
Yes	404	148(36.6%)	256(63.4%)		

*Chi-squared Test ( 2) was done to analyze the data. \*significant*

Table 2 shows some variables like daily duration of Internet usage ( $P<0.001^*$ ) and monthly expenditure for internet access ( $P=0.001^*$ ) were found significantly associated with Internet Addiction. Internet Addiction is relatively higher in the students who were always online ( $P<0.001^*$ ) in comparison to others. Students who did not reach the prescribed quantity of physical activity had a much greater rate of Internet addiction ( $P<0.001^*$ ). Among medical students those who maintained social distancing in the time of pandemic showed significant Association ( $P=0.008^*$ ) with Internet Addiction.

**Table 3. Multiple logistic regression for predictors of Internet Addiction of medical students in COVID-19 pandemic period**

Variables	$\beta$	S.E.	p-value	AOR	95% CI for OR	
					Lower	Upper
Lives with other people	1.257	.238	.000	3.515	2.203	5.607

Academic year (2nd)	.329	.260	.206	1.389	.835	2.312
Academic year (5th)	-0.767	.494	.120	0.464	.176	1.223
Daily time spend (> 4 hours)	.730	.265	.006	2.076	1.234	3.492
Monthly expenditure for internet access (>800 Tk.)	.549	.229	.016	1.732	1.106	2.713
Always online on social media sites	.844	.228	.000	2.326	1.487	3.639
Doesn't meets recommended PA threshold	.588	.234	.012	1.801	1.139	2.848
Believe in the past few months, your internet usage has increased due to maintain social distancing	1.139	.541	.035	3.123	1.081	9.025
Constant	-1.560	.295	.000	.210		

Table 3 presents the result from multiple logistic regression which tied students who lived with other people to Internet Addiction (AOR: 3.515, 95% CI: 2.203- 5.607,  $p = .000$ ). Within the criteria we defined, daily time spend (> 4 hours) (AOR: 2.076; 95% CI: 1.234- 3.492,  $p = 0.006$ ); monthly expenditure (>800Tk) (AOR: 1.732; 95% CI: 1.106-2.713,  $p = 0.016$ ); and always logging status in social medial (AOR: 2.326; 95% CI: 1.487-3.639,  $p = 0.000$ ) were the most significant predictors of medical students' Internet Addiction. Students who did not meet recommended physical activities threshold were more prone to be addicted (AOR: 1.801; 95% CI: 1.139 -2.848,  $p = 0.012$ ) in comparison to those who meet recommended physical activity. Furthermore, students who reported greater internet usage as a result of maintaining social distance were shown to be more likely to acquire Internet Addiction (AOR: 3.123, 95% CI: 1.081 - 9.025;  $p = 0.035$ ). The academic year of the students (2nd Year and 5th Year) failed to show any significant statistical association with respect to Internet Addiction in this regression model, but this one is significant in chi-square analysis.

## Discussion

In this survey, we discovered that 34.9 percent of people are addicted to the internet. Hassan et al. (2020) reported a prevalence of 27.1 percent among graduate students at Dhaka University in a study which was conducted in the three divisions of Bangladesh: Dhaka, Chittagong, and Sylhet, and Islam and Hossin (2016) reported a prevalence of nearly 24 percent among graduate students at Dhaka University in another study. The current findings, however, are significantly more prevalent among students than prior research, which can be related to the COVID-19. The research also revealed that students who stayed with others, as in a dormitory or a mess, have a higher possibility for developing an Internet addiction comparing to students living with their families. This results aligns with Hassan et al. (2020), who found that students who live with others have a higher Internet Addiction score. Loneliness, a lack of family bonding, online privacy, and a lack of monitoring and care are all possible contributing factors (Chaudhari et al., 2015; Uddin et al., 2016).

Internet addict students were shown to spend much more money and time on the Internet than non-addict students in this study. This conclusion is consistent with earlier research (Ghosh & Chatterjee, 2018; Chaudhari et al., 2015; Hassan et al., 2020). Internet addicts' increased spend of money and time may result in financial issues and a lack of focus on academic goals.

A significant statistical association between Internet Addiction and those who preferred to always stay online was present in the data. Nath et al. (2016), in their study among medical students in Northeastern India, reported that permanent login status highly associated with Internet Addiction.

Internet addiction was also shown to be more prevalent among those who did not engage in any physical exercise. A previous study focused on young adults and graduate students of Bangladesh also found physical inactivity as a risk factor for Internet Addiction (Hassan et al., 2020; Islam and Hossin, 2016). Furthermore, because to the COVID-19 epidemic, the maintaining of social distance has risen in recent months. However, unlike many other countries, Indonesia did not implement a mandatory "lockdown" throughout the nation during the COVID-19 epidemic, and social distance was not shown to be a risk of Internet Addiction among adult Indonesians (Siste et al., 2020)

Fifth-year students were shown to be more hooked than other students in our study. A recent study by Uddin et al. (2016) confirmed the findings, demonstrating that fourth-year undergraduate students are more hooked than those in prior years (1st, 2nd, and 3rd). However, this is due to excessive searching of study materials for their final year exam preparation, which makes them more prone to Internet. During this time, they may be more addicted to the Internet than other year medical students.

## Conclusion



The result from this study unfolds the severity and spread of Internet Addiction among medical students, and this requires timely remedial actions. It is imperative to provide anticipatory actions and fortify education on Internet Addiction among medical students while directing combined effort in mainstreaming mental health literacy. Strict observation and close monitoring are needed to overcome the long time use of Internet for recreational purposes. That there are a high possibility and risk of Internet Addiction should be conveyed with emphasis to the students, their guardians, and other concerned authorities. Awareness programs targeting all of the mentioned groups of people will ease the process of intervention and implementation of required policies at every levels of the society. Although Internet use for research and creative purposes should be encouraged and promoted, awareness about the demerits of uncontrolled and extended internet use should be spread among the young generation. Ultimately, we must be well acquainted with the advantage of the Internet, which should not be outweighed by the unhealthy consequences borne by its addiction.

## References

1. Bates, L. C., Zieff, G., Stanford, K., Moore, J. B., Kerr, Z. Y., Hanson, E. D., Gibbs, B. B., Kline, C. E., & Stoner, L. (2020). COVID-19 impact on behaviors across the 24-hour day in children and adolescents: physical activity, sedentary behavior, and sleep. *Children*, 7(9), 138.
2. Nath, K., Naskar, S., & Victor, R. (2016). A cross-sectional study on the prevalence, risk factors, and ill effects of internet addiction among medical students in Northeastern India. *The primary care companion for CNS disorders*, 18(2).
3. Ghosh, S., & Chatterjee, S. (2018). Assessment of Internet Addiction among Undergraduate Medical Students: A Cross-Sectional Study in a Medical College of Kolkata. *Journal of Clinical & Diagnostic Research*, 12(4).
4. Poli, R., & Agrimi, E. (2012). Internet addiction disorder: prevalence in an Italian student population. *Nordic journal of psychiatry*, 66(1), 55-59.
5. Uddin, M. S., Al Mamun, A., Iqbal, M. A., Nasrullah, M., Asaduzzaman, M., Sarwar, M. S., & Amran, M. S. (2016). Internet addiction disorder and its pathogenicity to psychological distress and depression among university students: A cross-sectional pilot study in Bangladesh. *Psychology*, 7(08), 1126.
6. Salicetia, F. (2015). Internet addiction disorder (IAD). *Procedia-Social and Behavioral Sciences*, 191, 1372-1376.
7. Haand, R., & Shuwang, Z. (2020). The relationship between social media addiction and depression: a quantitative study among university students in Khost, Afghanistan. *International Journal of Adolescence and Youth*, 25(1), 780-786.
8. Weinstein, A., & Lejoyeux, M. (2010). Internett-avhengighet eller overdreven bruk av Internett. *The American Journal of Drug and Alcohol Abuse*, 36(5), 277-283.
9. Mamun, M. A., Rafi, M. A., Al Mamun, A. S., Hasan, M. Z., Akter, K., Hsan, K., & Griffiths, M. D.


- (2019). Prevalence and psychiatric risk factors of excessive internet use among northern Bangladeshi job-seeking graduate students: a pilot study. *International Journal of Mental Health and Addiction*, 1-11.
10. Islam, M. A., & Hossain, M. Z. (2016). Prevalence and risk factors of problematic internet use and the associated psychological distress among graduate students of Bangladesh. *Asian Journal of Gambling Issues and Public Health*, 6(1), 1-14.
  11. Chaudhari, B., Menon, P., Saldanha, D., Tewari, A., & Bhattacharya, L. (2015). Internet addiction and its determinants among medical students. *Industrial psychiatry journal*, 24(2), 158.
  12. Hassan, T., Alam, M. M., Wahab, A., & Hawlader, M. D. (2020). Prevalence and associated factors of internet addiction among young adults in Bangladesh. *Journal of the Egyptian Public Health Association*, 95(1), 1-8.
  13. Karim, A. R., & Nigar, N. (2014). The internet addiction test: Assessing its psychometric properties in Bangladeshi culture. *Asian Journal of Psychiatry*, 10, 75-83.
  14. Mamun, M. A., & Griffiths, M. D. (2019). The assessment of internet addiction in Bangladesh: Why are prevalence rates so different? *Asian Journal of Psychiatry*.
  15. Hall, G., Laddu, D. R., Phillips, S. A., Lavie, C. J., & Arena, R. (2021). A tale of two pandemics: How will COVID-19 and global trends in physical inactivity and sedentary behavior affect one another?. *Progress in cardiovascular diseases*, 64, 108.
  16. Garcia-Priego, B. A., Triana-Romero, A., Pinto-Galvez, S. M., Duran-Ramos, C., Salas-Nolasco, O., Reyes, M. M., Ramos-de-la-Medina, A., & Troche, J. M. R. (2020). Anxiety, depression, attitudes, and internet addiction during the initial phase of the 2019 coronavirus disease (COVID-19) epidemic: A cross-sectional study in México. *MedRxiv*.
  17. Dong, H., Yang, F., Lu, X., & Hao, W. (2020). Internet addiction and related psychological factors among children and adolescents in China during the coronavirus disease 2019 (COVID-19) epidemic. *Frontiers in Psychiatry*, 11, 751.
  18. Ali, R., Mohammed, N., & Aly, H. (2017). Internet addiction among medical students of Sohag University, Egypt. *Journal of Egyptian Public Health Association*, 92(2), 86-95.
  19. Siste, K., Hanafi, E., Lee Thung Sen, H. C., Adrian, L. P. S., Limawan, A. P., Murtani, B. J., & Suwartono, C. (2020). The impact of physical distancing and associated factors towards internet addiction among adults in Indonesia during COVID-19 pandemic: a nationwide web-based study. *Frontiers in psychiatry*, 11.

## Author Bibliography

	<p><b>Nusrat Jahan</b> Dr. Nusrat Jahan has been teaching at Ibrahim Medical College's Department of Biochemistry in Dhaka, Bangladesh, for the past 9 years. She holds an MBBS degree and also has a Masters in Public Health degree. She is very interested in studying how social events interplay with the health of general population and how positive change in a country's public health can be encouraged. Email: <a href="mailto:nusrat27rumman@gmail.com">nusrat27rumman@gmail.com</a></p>
	<p><b>Md. Rumman Idris</b> Lt. Col. Dr. Rumman Idris have been working at the Cardiovascular Center in Combined Military Hospital (CMH), Dhaka, Bangladesh since 2015. He is an Assistant Professor and surgeon specializing in cardio-thoracic surgery at the center. Email: <a href="mailto:rummanidris2498@gmail.com">rummanidris2498@gmail.com</a></p>



---

	<p><b>Iffat Tania Prianka</b> Dr. Iffat Tania Prianka has joined as a lecturer in the Department of Biochemistry and later joined the Department of Community Medicine in Ibrahim Medical College. She is passionate to do Public health research and currently am pursuing M.phil in Preventive and social medicine from NIPSOM. Email: <a href="mailto:iffatprianka@gmail.com">iffatprianka@gmail.com</a></p>
	<p><b>Pradip Kumar Sen Gupta</b> Dr. Pradip Kumar Sen Gupta is Professor and currently is the head of the Department of Epidemiology of Bangladesh University of Health Science. He is also a faculty member of the Public Health Department at American International University, Bangladesh. Email: <a href="mailto:psen.06@gmail.com">psen.06@gmail.com</a></p>